

Finding the Holy Grail Is Not a Short-Term Project

Clare L. Atzema, MD, MSc; Michael J. Schull, MD, MSc

Clinical decision instruments exist for a myriad of disease states as they present to the emergency department (ED), including cervical spine imaging rules,^{1,2} instruments that estimate the risk of venous thromboembolism,³ and the mortality risk associated with pneumonia,⁴ to name a few. As the second most common reason to visit an ED,⁵ chest pain has been the target of decades of clinical decision instrument research efforts.

Article see p 195

Given that less than a quarter of these patients are suffering from an acute coronary syndrome,⁶ multiple groups have attempted to derive a highly sensitive decision instrument that identifies essentially all the patients with an acute coronary syndrome, while at the same time facilitating the safe, early discharge of a cohort of low-risk patients.⁷⁻¹⁰ Emergency physicians have identified 1% as the maximum miss rate that would be acceptable to them in order to actually use such a tool in the clinical setting.^{11,12} A tool that has a sensitivity with a lower confidence interval of 99%, but still has a high enough specificity to identify a substantial proportion of chest pain patients for ED discharge would be the holy grail of clinical decision instruments.

Early instruments had poor clinical uptake because of unacceptably low sensitivity: these include the Goldman Risk score, acute cardiac ischemia time-insensitive predictive instrument (ACI-TIPI), the Thrombolysis in Myocardial Infarction (TIMI) risk score, and Global Registry of Acute Coronary Events (GRACE).¹³⁻¹⁶ More recently, the North American Chest Pain Rule, the Accelerated Diagnostic Protocol to Assess Patients With Chest Pain Symptoms Using Contemporary Troponins (ADAPT) study, and the HEART (History, Electrocardiogram, Age, Risk factors, Troponin) Pathway, among others, have incorporated conventional troponins into their clinical decision instruments.^{7,8,10}

The North American Chest Pain Rule demonstrated a sensitivity of 100% (95% confidence interval, 97.2–100) in the derivation phase, and allowed for the early discharge of 18% of patients. Although the sensitivity remained high in subsequent

validation studies, the proportion of patients eligible for early discharge dropped to 11%.¹⁷ The Asia-Pacific Evaluation of Chest pain Trial (ASPECT) study¹² also had high sensitivity, but only 10% were identified for early discharge, although the subsequent ADAPT tool¹⁰ identified 20% for discharge. The latter tool has been validated using high-sensitivity troponins, and maintained the high sensitivity while identifying >20% of patients for early ED discharge.¹⁸ The HEART score has evolved into the HEART Pathway, which showed 100% sensitivity using 2 sets of conventional troponins, although it had a lower confidence interval of 72%.⁸ An external validation study of the HEART Pathway had a miss rate of 1.7% (95% confidence interval, 1.0–2.9); lowering the low-risk score cut-off from 3 to 2 resulted in a miss rate 1.1%.¹⁹ It is clear that researchers are getting closer to a clinically acceptable tool for the safe early disposition of patients with chest pain from ED, but no instrument seems ready for prime-time yet.

In this issue of *Circulation: Cardiovascular Quality and Outcomes*, Mahler et al²⁰ assessed objective cardiac testing in ED patients with chest pain up to 30 days after the ED visit, using a rigorously conducted randomized control trial. Patients were randomized to the HEART Pathway (where a HEART score of ≤ 3 was used to define low risk) or standard of care. In addition to objective cardiac testing, the authors compared time spent in the ED and hospital, the proportion of patients who achieved early ED discharge (discharge without objective cardiac testing), and major adverse cardiac events (MACE) at 30 days. The study found a substantial reduction in objective cardiac testing up to 30 days after the ED visit among the HEART Pathway patients, relative to controls. In addition, patients in the HEART pathway arm had substantially shorter times in the ED and hospital than control patients, and more were discharged early from the ED (21% increase in early discharge compared with controls, compared with 8% more using the ADAPT tool²¹).

There was no difference in MACE at 30 days; however, it is important to note that the study was not powered to find a difference in MACE. Although one might look to the previously published, observational HEART studies (that showed high sensitivities for MACE) to confirm the safety of this approach, post-ED diagnostic testing and interventions that occur in observational studies may have decreased the rate of 30-day MACE. In other words, more episodes of MACE might have occurred if further testing had not been performed. In addition, although the authors present this as a real-world test of the HEART Pathway, the clinicians in this RCT were alerted if they had miscalculated the patient's HEART score (underestimated it), which may also have led to underestimation of MACE. These issues will need to be addressed prior to clinical use of the HEART Pathway.

A similar RCT has been conducted for the ADAPT tool²¹: that study included comparable outcomes, with the exception

From the Institute for Clinical Evaluative Sciences, Toronto, Ontario, Canada (C.L.A., M.J.S.); Division of Emergency Medicine, Department of Medicine, University of Toronto, Toronto, Ontario, Canada (C.L.A., M.J.S.); and Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada (C.L.A., M.J.S.).

The opinions expressed in this article are not necessarily those of the editors or of the American Heart Association.

Correspondence to Clare Atzema, MD, MSc, 2075 Bayview Ave, G146, Toronto, ON M4N 3M5, Canada. E-mail clare.atzema@ices.on.ca (*Circ Cardiovasc Qual Outcomes*. 2015;8:135-137.

DOI: 10.1161/CIRCOUTCOMES.115.001611.)

© 2015 American Heart Association, Inc.

Circ Cardiovasc Qual Outcomes is available at <http://circoutcomes.ahajournals.org>

DOI: 10.1161/CIRCOUTCOMES.115.001611

of objective cardiac testing up to 30 days after the ED visit. The authors of the ADAPT studies have made it clear that patients discharged using their accelerated diagnostic protocol still require further nonurgent follow-up investigations.^{10,18} This is because of the aforementioned limitation of observational studies, where subsequent investigations and treatments may have lowered the frequency of MACE. In comparison, the study by Mahler et al²⁰ takes the goal of getting low-risk chest pain patients out of the ED and hospital one step further, by assessing *both* inpatient and outpatient objective cardiac testing up to 30 days after the ED visit, with the aim of reducing it as well (in comparison, in the ADAPT study all patients received objective cardiac testing within 72 hours of ED discharge).²¹

Clinically, the leap of faith required for this outcome measure is sizeable. Even in Canada, where physicians are far less likely to request objective testing before ED discharge,^{7,9} we suspect that removing the safeguard of subsequent outpatient objective cardiac testing on patients with a reasonable presentation for acute coronary syndrome is not likely to gain widespread traction. In one Canadian study, this is how 17% of the acute coronary syndromes were identified.⁹

Therefore, although the study by Mahler et al²⁰ showed a decrease in objective cardiac testing up to 30 days after ED discharge, the key question to ask is whether that is a good thing. The American College of Cardiology/American Heart Association recommends that patients deemed low risk in the ED setting should undergo objective cardiac testing, either in hospital, in an ED observation unit, or in the early period after ED discharge.²² However, there are no studies with long-term outcomes to substantiate, or disprove, this one-size-fits-all approach. Most emergency physicians are aware that there is a high rate of false positives associated with objective testing of low-risk patients, and would like to avoid sending such patients for testing, consistent with the Choosing Wisely campaign.²³ Specifically, what they want to reduce is *inappropriate* objective cardiac testing in low-risk chest pain patients, as opposed to reducing outpatient cardiac testing overall.

A valuable finding in the study by Mahler et al²⁰ was that a substantial minority (29%) of discharged low-risk patients in the HEART Pathway arm received subsequent objective cardiac testing within 30 days anyway. The rate of testing would probably have been higher had follow-up been longer than 30 days, raising the question of whether the HEART Pathway simply delays objective testing as opposed to forgoing it. The family physicians and outpatient specialists who took over the care of these patients after ED discharge apparently did not feel comfortable trusting in the decision instrument's recommendations; this finding highlights the need to engage the practitioners who are tasked with the long-term care of these patients, to effectively implement such a decision instrument in clinical care. Finally, it would be interesting to incorporate the patient's perspective on the risks and benefits associated with the decision instrument's recommendations, as well as on a care pathway approach that focuses on what the patient does not have, as opposed to what is causing their chest pain.

The study by Mahler et al²⁰ is to be applauded for highlighting the understudied nature of the American College of Cardiology/American Heart Association guideline

recommendations for objective cardiac testing. We look forward to future studies by this group that are powered to determine the short- and long-term effect of the HEART Pathway on outcomes of ED chest pain patients. Successful clinical implementation of a highly sensitive tool that can facilitate the safe early discharge of more ED chest pain patients would be a tremendous gain for healthcare systems around the world, and defining a group of patients who do not need outpatient objective cardiac testing after ED discharge would be a further coup.

Sources of Funding

Dr Atzema is funded by a New Investigator Award from the Heart and Stroke Foundation of Ontario, and Dr Schull is supported by an Applied Chair in Health Services and Policy Research from CIHR.

Disclosures

None.

References

- Hoffman JR, Mower WR, Wolfson AB, Todd KH, Zucker MI. Validity of a set of clinical criteria to rule out injury to the cervical spine in patients with blunt trauma. National Emergency X-Radiography Utilization Study Group. *N Engl J Med*. 2000;343:94–99. doi: 10.1056/NEJM200007133430203.
- Stiell IG, Wells GA, Vandemheen KL, Clement CM, Lesiuk H, De Maio VJ, Laupacis A, Schull M, McKnight RD, Verbeek R, Brison R, Cass D, Dreyer J, Eisenhauer MA, Greenberg GH, MacPhail I, Morrison L, Reardon M, Worthington J. The Canadian C-spine rule for radiography in alert and stable trauma patients. *JAMA*. 2001;286:1841–1848.
- Wells PS, Anderson DR, Rodger M, Stiell I, Dreyer JF, Barnes D, Forgie M, Kovacs G, Ward J, Kovacs MJ. Excluding pulmonary embolism at the bedside without diagnostic imaging: management of patients with suspected pulmonary embolism presenting to the emergency department by using a simple clinical model and d-dimer. *Ann Intern Med*. 2001;135:98–107.
- Fine MJ, Auble TE, Yealy DM, Hanusa BH, Weissfeld LA, Singer DE, Coley CM, Marrie TJ, Kapoor WN. A prediction rule to identify low-risk patients with community-acquired pneumonia. *N Engl J Med*. 1997;336:243–250. doi: 10.1056/NEJM199701233360402.
- Niska R, Bhuiya F, Xu J. National Hospital Ambulatory Medical Care Survey: 2007 emergency department summary. *Natl Health Stat Report*. 2010;1–31.
- Pope JH, Aufderheide TP, Ruthazer R, Woolard RH, Feldman JA, Beshansky JR, Griffith JL, Selker HP. Missed diagnoses of acute cardiac ischemia in the emergency department. *N Engl J Med*. 2000;342:1163–1170. doi: 10.1056/NEJM200004203421603.
- Hess EP, Brison RJ, Perry JJ, Calder LA, Thiruganasambandamoorthy V, Agarwal D, Sadosty AT, Silvillotti ML, Jaffe AS, Montori VM, Wells GA, Stiell IG. Development of a clinical prediction rule for 30-day cardiac events in emergency department patients with chest pain and possible acute coronary syndrome. *Ann Emerg Med*. 2012;59:115–125. doi: 10.1016/j.annemergmed.2011.07.026.
- Mahler SA, Hiestand BC, Goff DC Jr, Hoekstra JW, Miller CD. Can the HEART score safely reduce stress testing and cardiac imaging in patients at low risk for major adverse cardiac events? *Crit Pathw Cardiol*. 2011;10:128–133. doi: 10.1097/HPC.0b013e3182315a85.
- Scheuermeyer FX, Innes G, Grafstein E, Kiess M, Boychuk B, Yu E, Kalla D, Christenson J. Safety and efficiency of a chest pain diagnostic algorithm with selective outpatient stress testing for emergency department patients with potential ischemic chest pain. *Ann Emerg Med*. 2012;59:256–264. doi: 10.1016/j.annemergmed.2011.10.016.
- Than M, Cullen L, Aldous S, Parsonage WA, Reid CM, Greenslade J, Flaws D, Hammett CJ, Beam DM, Ardagh MW, Troughton R, Brown AF, George P, Florkowski CM, Kline JA, Peacock WF, Maisel AS, Lim SH, Lamanna A, Richards AM. 2-Hour accelerated diagnostic protocol to assess patients with chest pain symptoms using contemporary troponins as the only biomarker: the ADAPT trial. *J Am Coll Cardiol*. 2012;59:2091–2098. doi: 10.1016/j.jacc.2012.02.035.
- Kline JA, Johnson CL, Pollack CV Jr, Diercks DB, Hollander JE, Newgard CD, Garvey JL. Pretest probability assessment derived from attribute matching. *BMC Med Inform Decis Mak*. 2005;5:26. doi: 10.1186/1472-6947-5-26.

12. Than M, Cullen L, Reid CM, Lim SH, Aldous S, Ardagh MW, Peacock WF, Parsonage WA, Ho HF, Ko HF, Kasliwal RR, Bansal M, Soerianata S, Hu D, Ding R, Hua Q, Seok-Min K, Sritara P, Sae-Lee R, Chiu TF, Tsai KC, Chu FY, Chen WK, Chang WH, Flaws DF, George PM, Richards AM. A 2-h diagnostic protocol to assess patients with chest pain symptoms in the Asia-Pacific region (ASPECT): a prospective observational validation study. *Lancet*. 2011;377:1077–1084. doi: 10.1016/S0140-6736(11)60310-3.
13. de Araújo Gonçalves P, Ferreira J, Aguiar C, Seabra-Gomes R. TIMI, PURSUIT, and GRACE risk scores: sustained prognostic value and interaction with revascularization in NSTE-ACS. *Eur Heart J*. 2005;26:865–872. doi: 10.1093/eurheartj/ehi187.
14. Goldman L, Cook EF, Brand DA, Lee TH, Rouan GW, Weisberg MC, Acampora D, Stasiulewicz C, Walshon J, Terranova G. A computer protocol to predict myocardial infarction in emergency department patients with chest pain. *N Engl J Med*. 1988;318:797–803. doi: 10.1056/NEJM198803313181301.
15. Lyon R, Morris AC, Caesar D, Gray S, Gray A. Chest pain presenting to the Emergency Department—to stratify risk with GRACE or TIMI? *Resuscitation*. 2007;74:90–93. doi: 10.1016/j.resuscitation.2006.11.023.
16. Selker HP, Beshansky JR, Griffith JL, Aufderheide TP, Ballin DS, Bernard SA, Crespo SG, Feldman JA, Fish SS, Gibler WB, Kiez DA, McNutt RA, Moulton AW, Ornato JP, Podrid PJ, Pope JH, Salem DN, Sayre MR, Woolard RH. Use of the acute cardiac ischemia time-insensitive predictive instrument (ACI-TIPI) to assist with triage of patients with chest pain or other symptoms suggestive of acute cardiac ischemia. A multicenter, controlled clinical trial. *Ann Intern Med*. 1998;129:845–855.
17. Mahler SA, Miller CD, Hollander JE, Nagurney JT, Birkhahn R, Singer AJ, Shapiro NI, Glynn T, Nowak R, Safdar B, Peberdy M, Counselman FL, Chandra A, Kosowsky J, Neuenschwander J, Schrock JW, Plantholt S, Diercks DB, Peacock WF. Identifying patients for early discharge: performance of decision rules among patients with acute chest pain. *Int J Cardiol*. 2013;168:795–802. doi: 10.1016/j.ijcard.2012.10.010.
18. Cullen L, Mueller C, Parsonage WA, Wildi K, Greenslade JH, Twerenbold R, Aldous S, Meller B, Tate JR, Reichlin T, Hammett CJ, Zellweger C, Ungerer JP, Rubini Gimenez M, Troughton R, Murray K, Brown AF, Mueller M, George P, Mosimann T, Flaws DF, Reiter M, Lamanna A, Haaf P, Pemberton CJ, Richards AM, Chu K, Reid CM, Peacock WF, Jaffe AS, Florkowski C, Deely JM, Than M. Validation of high-sensitivity troponin I in a 2-hour diagnostic strategy to assess 30-day outcomes in emergency department patients with possible acute coronary syndrome. *J Am Coll Cardiol*. 2013;62:1242–1249. doi: 10.1016/j.jacc.2013.02.078.
19. Six AJ, Cullen L, Backus BE, Greenslade J, Parsonage W, Aldous S, Doevendans PA, Than M. The HEART score for the assessment of patients with chest pain in the emergency department: a multinational validation study. *Crit Pathw Cardiol*. 2013;12:121–126. doi: 10.1097/HPC.0b013e31828b327e.
20. Mahler SA, Riley RF, Hiestand BC, Russell GB, Hoekstra JW, Lefebvre CW, Nicks BA, Cline DM, Askew KL, Elliott SB, Herrington DM, Burke GL, Miller CD. The HEART Pathway randomized trial: identifying emergency department patients with acute chest pain for early discharge. *Circ Cardiovasc Qual Outcomes*. 2015; 8: 195–203. doi: 10.1161/CIRCOUTCOMES.114.001384.
21. Than M, Aldous S, Lord SJ, Goodacre S, Frampton CM, Troughton R, George P, Florkowski CM, Ardagh M, Smyth D, Jardine DL, Peacock WF, Young J, Hamilton G, Deely JM, Cullen L, Richards AM. A 2-hour diagnostic protocol for possible cardiac chest pain in the emergency department: a randomized clinical trial. *JAMA Intern Med*. 2014;174:51–58. doi: 10.1001/jamainternmed.2013.11362.
22. Anderson JL, Adams CD, Antman EM, Bridges CR, Califf RM, Casey DE Jr, Chavey WE II, Fesmire FM, Hochman JS, Levin TN, Lincoff AM, Peterson ED, Theroux P, Wenger NK, Wright RS, Smith SC Jr, Jacobs AK, Halperin JL, Hunt SA, Krumholz HM, Kushner FG, Lytle BW, Nishimura R, Ornato JP, Page RL, Riegel B; American College of Cardiology; American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non ST-Elevation Myocardial Infarction); American College of Emergency Physicians; Society for Cardiovascular Angiography and Interventions; Society of Thoracic Surgeons; American Association of Cardiovascular and Pulmonary Rehabilitation; Society for Academic Emergency Medicine. ACC/AHA 2007 guidelines for the management of patients with unstable angina/non ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non ST-Elevation Myocardial Infarction): developed in collaboration with the American College of Emergency Physicians, the Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons; endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation and the Society for Academic Emergency Medicine. *Circulation*. 2007;116:e148–e304. doi: 10.1161/CIRCULATIONAHA.107.181940.
23. ABIM Foundation. Choosing wisely. <http://www.choosingwisely.org/resources/2015>. Accessed January 5, 2015.

KEY WORDS: Editorials ■ acute coronary syndrome ■ decision support techniques ■ hospital emergency service ■ patient discharge

Finding the Holy Grail Is Not a Short-Term Project

Clare L. Atzema and Michael J. Schull

Circ Cardiovasc Qual Outcomes. 2015;8:135-137; originally published online March 3, 2015;
doi: 10.1161/CIRCOUTCOMES.115.001611

Circulation: Cardiovascular Quality and Outcomes is published by the American Heart Association, 7272
Greenville Avenue, Dallas, TX 75231

Copyright © 2015 American Heart Association, Inc. All rights reserved.

Print ISSN: 1941-7705. Online ISSN: 1941-7713

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://circoutcomes.ahajournals.org/content/8/2/135>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation: Cardiovascular Quality and Outcomes* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation: Cardiovascular Quality and Outcomes* is online at:
<http://circoutcomes.ahajournals.org//subscriptions/>